

**Interesting quantum states
(Lecture of the Quantum Information class of
the Master in Quantum Science and
Technology)**

Géza Tóth

Theoretical Physics, University of the Basque Country (UPV/EHU), Bilbao, Spain
Donostia International Physics Center (DIPC), San Sebastián, Spain
IKERBASQUE, Basque Foundation for Science, Bilbao, Spain
Wigner Research Centre for Physics, Budapest, Hungary

UPV/EHU, Leioa
22, 23 January, 2024

1 Interesting quantum states

- Motivation
- A. Single particle states
- B. Bipartite singlet state
- C. Werner states
- D. Isotropic states
- E. Schrödinger cat states
- F. Greenberger-Horne-Zeilinger (GHZ) state
- G. W state
- H. Symmetric Dicke states

Which quantum states are interesting?

- We have infinite possibilities to pick a quantum state in a multi-qubit system.
- We would like to find useful ones or states that have interesting symmetries.

1 Interesting quantum states

- Motivation
- A. Single particle states
- B. Bipartite singlet state
- C. Werner states
- D. Isotropic states
- E. Schrödinger cat states
- F. Greenberger-Horne-Zeilinger (GHZ) state
- G. W state
- H. Symmetric Dicke states

Single particle states

- Pure states. The von Neumann entropy $S = 0$.
- Completely mixed state

$$\rho_{\text{cm}} = \frac{1}{d} \sum_{k=1}^d |k\rangle\langle k|.$$

The von Neumann entropy $S = \log d$, maximal.

1 Interesting quantum states

- Motivation
- A. Single particle states
- **B. Bipartite singlet state**
- C. Werner states
- D. Isotropic states
- E. Schrödinger cat states
- F. Greenberger-Horne-Zeilinger (GHZ) state
- G. W state
- H. Symmetric Dicke states

Bipartite singlet state

- The two-qubit singlet state looks like

$$|\Psi_{\text{singlet}}\rangle = \frac{1}{\sqrt{2}} (|01\rangle - |10\rangle).$$

- We get the same form after any basis transformation (if we transform the bases of the two qubits in the same way). This can be seen as follows. Let us choose two vectors as

$$\begin{aligned} |v\rangle &= \alpha|0\rangle + \beta|1\rangle, \\ |v_{\perp}\rangle &= \beta^*|0\rangle - \alpha^*|1\rangle. \end{aligned}$$

Clearly,

$$\langle v|v_{\perp}\rangle = 0,$$

Then, simple algebra yields

$$\frac{1}{\sqrt{2}} (|v\rangle \otimes |v_{\perp}\rangle - |v_{\perp}\rangle \otimes |v\rangle) = \frac{1}{\sqrt{2}} (|01\rangle - |10\rangle).$$

This is true for any $|v\rangle$ and $|v_{\perp}\rangle$.

Bipartite singlet state II

- Due to the independence from the choice of the local basis, it is invariant under a transformation of the type $U \otimes U$, apart from a global phase ϕ .

$$U \otimes U |\Psi_{\text{singlet}}\rangle = |\Psi_{\text{singlet}}\rangle \exp(-i\phi).$$

We can also say that

$$U \otimes U |\Psi_{\text{singlet}}\rangle \langle \Psi_{\text{singlet}}| (U \otimes U)^\dagger = |\Psi_{\text{singlet}}\rangle \langle \Psi_{\text{singlet}}|.$$

Hence,

$$U \otimes U |\Psi_{\text{singlet}}\rangle \langle \Psi_{\text{singlet}}| = |\Psi_{\text{singlet}}\rangle \langle \Psi_{\text{singlet}}| U \otimes U.$$

Thus, the density matrices of such states will commute with all $U \otimes U$:

$$[U \otimes U, |\Psi_{\text{singlet}}\rangle \langle \Psi_{\text{singlet}}|] = 0 \quad (1)$$

for any U .

Bipartite singlet state III

- Let us consider some operators of the form

$$\sigma_{\vec{n}} = \sum_{l=x,y,z} n_l \sigma_l$$

where $|\vec{n}| = 1$. For $\vec{n} = (1, 0, 0)$, $\sigma_{\vec{n}} = \sigma_x$. For $\vec{n} = (0, 1, 0)$, $\sigma_{\vec{n}} = \sigma_y$, and in general it is a generalization of the Pauli spin matrices to an arbitrary direction.

- Such operators all have eigenvalues ± 1 . If you measure $\sigma_{\vec{n}}$ on party A and get a result, then if you also measure it on party B , you will get the opposite result. This is true for every $\sigma_{\vec{n}}$.
- This can be used in quantum communication to establish a bit sequence that is known only by Alice and Bob and by nobody else.

Bipartite singlet state IV

- For the singlet state

$$|\Psi_{\text{singlet}}\rangle = \frac{1}{\sqrt{2}} (|01\rangle - |10\rangle).$$

we have anticorrelations

$$\begin{aligned}\langle \sigma_x \otimes \sigma_x \rangle &= -1, \\ \langle \sigma_y \otimes \sigma_y \rangle &= -1, \\ \langle \sigma_z \otimes \sigma_z \rangle &= -1.\end{aligned}\tag{2}$$

- For the collective angular momentum, we have

$$[\Delta(\sigma_x^{(1)} + \sigma_x^{(2)})]^2 = [\Delta(\sigma_y^{(1)} + \sigma_y^{(2)})]^2 = [\Delta(\sigma_z^{(1)} + \sigma_z^{(2)})]^2 = 0.\tag{3}$$

Bipartite singlet state V

- Why is it called a singlet? Remember the theory of angular momentum, triplet and singlet subspace.
- Alternatively, in quantum information, the maximally entangled state

$$|\Psi_{\text{me}}\rangle = \frac{1}{\sqrt{2}} (|00\rangle + |11\rangle)$$

can also be called singlet.

- A generalization for higher dimensions is the maximally entangled state

$$|\Psi_{\text{me}}\rangle = \frac{1}{\sqrt{d}} \sum_{k=1}^d |kk\rangle.$$

- For the maximally entangled state, the reduced state is the completely mixed state

$$\text{Tr}_A(|\Psi_{\text{me}}\rangle\langle\Psi_{\text{me}}|) = \text{Tr}_B(|\Psi_{\text{me}}\rangle\langle\Psi_{\text{me}}|) = \frac{\mathbb{1}}{d}.$$

Thus, if we have access only to one of the two subsystems, we know nothing.

Bipartite singlet state VI

- For the maximally entangled state

$$|\Psi_{\text{me}}\rangle = \frac{1}{\sqrt{2}} (|00\rangle + |11\rangle)$$

we have

$$\begin{aligned}\langle\sigma_x \otimes \sigma_x\rangle &= +1, \\ \langle\sigma_y \otimes \sigma_y\rangle &= -1, \\ \langle\sigma_z \otimes \sigma_z\rangle &= +1.\end{aligned}\tag{4}$$

- For the collective angular momentum, we have

$$[\Delta(\sigma_x^{(1)} - \sigma_x^{(2)})]^2 = [\Delta(\sigma_y^{(1)} + \sigma_y^{(2)})]^2 = [\Delta(\sigma_z^{(1)} - \sigma_z^{(2)})]^2 = 0. \tag{5}$$

1 Interesting quantum states

- Motivation
- A. Single particle states
- B. Bipartite singlet state
- **C. Werner states**
- D. Isotropic states
- E. Schrödinger cat states
- F. Greenberger-Horne-Zeilinger (GHZ) state
- G. W state
- H. Symmetric Dicke states

Flip operator

- Flip operator

$$F|\Psi\rangle|\Phi\rangle = |\Phi\rangle|\Psi\rangle.$$

- For two-qubits

$$F = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

$$F = \frac{1}{2}(\mathbb{1} \otimes \mathbb{1} + \sigma_x \otimes \sigma_x + \sigma_y \otimes \sigma_y + \sigma_z \otimes \sigma_z).$$

- Eigenvalues of F : -1,1,1,1 (-1 for antisymmetric states, +1 for symmetric states).

Flip operator II

- For a comparison, remember that for the projector to the singlet state we have

$$|\Psi_{\text{singlet}}\rangle\langle\Psi_{\text{singlet}}| = \frac{1}{4}(\mathbb{1} \otimes \mathbb{1} - \sigma_x \otimes \sigma_x - \sigma_y \otimes \sigma_y - \sigma_z \otimes \sigma_z).$$

- For the projector to the maximally entangled state we have

$$|\Psi_{\text{me}}\rangle\langle\Psi_{\text{me}}| = \frac{1}{4}(\mathbb{1} \otimes \mathbb{1} + \sigma_x \otimes \sigma_x - \sigma_y \otimes \sigma_y + \sigma_z \otimes \sigma_z).$$

Werner states

- **Definition.** Werner states are states that are invariant under a transformation of the type $U \otimes U$ (Werner, 1989).
- For qubits, Werner states are noisy singlets for two-qubits

$$\rho_{\text{Werner}}(\rho) = (1 - \rho)|\Psi_{\text{singlet}}\rangle\langle\Psi_{\text{singlet}}| + \rho\frac{\mathbb{1}}{4}.$$

- For two qudits, their density matrix is defined as

$$\alpha\mathbb{1} + \beta F$$

where F is the flip operator. For systems larger than qubits, we do not have a pure Werner state.

- Twirling

$$\tau(\rho) = \int M(dU)U \otimes U \rho (U \otimes U)^\dagger.$$

Used to transform states into a normal form, for example, before distilling entanglement. Twirling leaves Werner states unchanged. It transforms all quantum states to Werner states.

1 Interesting quantum states

- Motivation
- A. Single particle states
- B. Bipartite singlet state
- C. Werner states
- **D. Isotropic states**
- E. Schrödinger cat states
- F. Greenberger-Horne-Zeilinger (GHZ) state
- G. W state
- H. Symmetric Dicke states

Isotropic states

- Isotropic states are invariant under any transformation of the type $U \otimes U^*$.
- Isotropic states are defined as the maximally entangled state

$$|\Psi_{\text{me}}\rangle = \frac{1}{\sqrt{d}} \sum_{k=1}^d |kk\rangle.$$

mixed with white noise.

- The maximally entangled state above is the pure isotropic state for any dimension.

1 Interesting quantum states

- Motivation
- A. Single particle states
- B. Bipartite singlet state
- C. Werner states
- D. Isotropic states
- **E. Schrödinger cat states**
- F. Greenberger-Horne-Zeilinger (GHZ) state
- G. W state
- H. Symmetric Dicke states

Schrödinger cat states

- Think on the usual the Schrödinger's cat experiment. The cat is in a superposition of being dead and alive.
- Questions about the linearity of quantum mechanics, etc. Can superpositions of macroscopically different objects exist?
- Funny thought-experiment: Wigner's friend.

1 Interesting quantum states

- Motivation
- A. Single particle states
- B. Bipartite singlet state
- C. Werner states
- D. Isotropic states
- E. Schrödinger cat states
- F. Greenberger-Horne-Zeilinger (GHZ) state
- G. W state
- H. Symmetric Dicke states

Greenberger-Horne-Zeilinger (GHZ) state

- A possible generalization of the maximally entangled state to N qubits is the GHZ state defined as

$$|\text{GHZ}\rangle = \frac{1}{\sqrt{2}} (|00\dots 00\rangle + |11\dots 11\rangle)$$

- The entanglement of the GHZ state is very fragile. Measuring one qubit destroys it, the state becomes separable:

$$\text{Tr}_1(|\text{GHZ}_N\rangle\langle\text{GHZ}_N|) = \frac{1}{2}(|0\rangle\langle 0|^{\otimes(N-1)} + |1\rangle\langle 1|^{\otimes(N-1)}).$$

- Realized experimentally with trapped ions, photons $\sim 10 - 15$ qubits.

Greenberger-Horne-Zeilinger (GHZ) state II

- Interestingly, the GHZ state is an eigenstate of operators that are the products of single-qubit operators. For example, it is the eigenstate of

$$\sigma_x \otimes \sigma_x \otimes \dots \otimes \sigma_x = \sigma_x^{\otimes N}$$

with an eigenvalue $+1$. What does this mean? If they flip all the qubits, we get back the original states.

- The GHZ state is also the eigenstate of the operators of the type

$$\sigma_z^{(m)} \sigma_z^{(n)},$$

for all $m \neq n$ with an eigenvalue $+1$.

Greenberger-Horne-Zeilinger (GHZ) state III

- If a state is eigenstate of operators O_1 and O_2 , then it is also an eigenstate of $O_1 O_2$. Because of that the state is an eigenstate of the products of such operators.
- Note that all these operators commute with each other.
- For all operators, we have $O^2 = \mathbb{1}$.
- These operators form a *group* that is called *stabilizer*. The group has N generators and 2^N elements.
- (Definition of a discrete group: If A and B are in the group, so is their product AB .)
- Stabilizer theory is important for quantum error correction.

Greenberger-Horne-Zeilinger (GHZ) state IV

- Just an example: for three-qubit GHZ states, we have the following 8 operators

$$\sigma_x \otimes \sigma_x \otimes \sigma_x,$$

$$\sigma_z \otimes \sigma_z \otimes \mathbb{1},$$

$$\mathbb{1} \otimes \sigma_z \otimes \sigma_z,$$

$$\sigma_z \otimes \mathbb{1} \otimes \sigma_z,$$

$$-\sigma_y \otimes \sigma_y \otimes \sigma_x,$$

$$-\sigma_x \otimes \sigma_y \otimes \sigma_y,$$

$$-\sigma_y \otimes \sigma_x \otimes \sigma_y,$$

$$\mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1}.$$

1 Interesting quantum states

- Motivation
- A. Single particle states
- B. Bipartite singlet state
- C. Werner states
- D. Isotropic states
- E. Schrödinger cat states
- F. Greenberger-Horne-Zeilinger (GHZ) state
- **G. W state**
- H. Symmetric Dicke states

W state

- It is defined as

$$\frac{1}{\sqrt{N}} (|100\dots\rangle + |010\dots\rangle + |001\dots\rangle + \dots + |000\dots01\rangle).$$

Interestingly, it maximizes the two-body concurrence among symmetric states.

- If we lose a single particle, we obtain

$$\text{Tr}_1(|W_N\rangle\langle W_N|) = \frac{1}{N}|0\rangle\langle 0|^{\otimes(N-1)} + \frac{N-1}{N}|W_{N-1}\rangle\langle W_{N-1}|.$$

The entanglement of the W state is robust. Losing one qubit does not destroy the entanglement.

- Realized with trapped ions, photons, cold atoms, up to thousands of particles.

1 Interesting quantum states

- Motivation
- A. Single particle states
- B. Bipartite singlet state
- C. Werner states
- D. Isotropic states
- E. Schrödinger cat states
- F. Greenberger-Horne-Zeilinger (GHZ) state
- G. W state
- H. Symmetric Dicke states

Symmetric Dicke states of qubits

PHYSICAL REVIEW

VOLUME 93, NUMBER 1

JANUARY 1, 1954

Coherence in Spontaneous Radiation Processes

R. H. DICKE

Palmer Physical Laboratory, Princeton University, Princeton, New Jersey

(Received August 25, 1953)

By considering a radiating gas as a single quantum-mechanical system, energy levels corresponding to certain correlations between individual molecules are described. Spontaneous emission of radiation in a transition between two such levels leads to the emission of coherent radiation. The discussion is limited first to a gas of dimension small compared with a wavelength. Spontaneous radiation rates and natural line breadths are calculated. For a gas of large extent the effect of photon recoil momentum on coherence is calculated. The effect of a radiation pulse in exciting "super-radiant" states is discussed. The angular correlation between successive photons spontaneously emitted by a gas initially in thermal equilibrium is calculated.

IN the usual treatment of spontaneous radiation by a gas, the radiation process is calculated as though the separate molecules radiate independently of each other. To justify this assumption it might be argued that, as a result of the large distance between molecules and subsequent weak interactions, the probability of a given molecule emitting a photon should be independent of the states of other molecules. It is clear that this model is incapable of describing a coherent spontaneous radiation process since the radiation rate is proportional to the molecular concentration rather than to the square of the concentration. This simplified picture overlooks the fact that all the molecules are interacting with a common radiation field and hence cannot be treated as independent. The model is wrong in principle and many of the results obtained from it are incorrect. A simple example will be used to illustrate the inade-

quacy of the usual treatment. Consider a system of two neutrons, each of which can be in a triplet or singlet state. The triplet state is capable of radiating to the ground state (triplet) but the singlet state will not couple with the triplet system. Consequently, only the triplet part is modified by the coupling with the field. After a long time there is still a probability of one-half that a photon has not been emitted. If, after a long period of time, no photon has been emitted, the neutrons are in a singlet state and it is impossible to predict which neutron is the excited one.

On the other hand, if the initial state of the two neutrons were triplet with $s=1$, $m_s=0$ namely a state with one excited neutron, a photon would be certain to be emitted and the transition probability would be just double that for a lone excited neutron. Thus, the presence of the unexcited neutron in this case doubles the radiation rate.

Symmetric Dicke states of qubits II

- Dicke states are simultaneous eigenstate of $\vec{J}^2 = J_x^2 + J_y^2 + J_z^2$ and J_z .
- Symmetric Dicke states of qubits have a maximal \vec{J}^2 , i.e.,

$$\langle J_x^2 + J_y^2 + J_z^2 \rangle = (N/2)(N/2 + 1). \quad (6)$$

They are the equal superpositions of the permutations of a series of 0's and 1's

$$|D_N^{(m)}\rangle \propto \sum_k \mathcal{P}_k(|0\rangle^{\otimes(N-m)} \otimes |1\rangle^{\otimes(m)}),$$

where the sum is over all distinct permutations of 0's and 1's.

Symmetric Dicke states of qubits III

- For $|0\rangle^{\otimes N}$, we have $\langle J_z \rangle = +N/2$.
- For $|1\rangle^{\otimes N}$, we have $\langle J_z \rangle = -N/2$.
- If half of the particles are 1, half of them are 0, we have $m = N/2$. In this case, $\langle J_z \rangle = 0$.

- For example, for $N = 4$ the symmetric Dicke state is

$$|D_4^{(2)}\rangle = \frac{1}{\sqrt{6}}(|0011\rangle + |0101\rangle + |1001\rangle + |0110\rangle + |1010\rangle + |1100\rangle).$$

- Relation to W states: $|D_N^{(1)}\rangle = |W_N\rangle$.
- Realized with trapped ions, photons, cold gases, up to thousands of particles.